

The s-wave implosion and explosion of free particles

I. Białynicki-Birula^{1,2}, M. A. Cirone², J. P. Dahl^{2,3}, M. Fedorov^{2,4}, W. P. Schleich²

¹ Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668, Warsaw, Poland

² Abteilung für Quantenphysik, Universität Ulm, D-89069 Ulm, Germany

³ Chemical Physics, Department of Chemistry, Technical University of Denmark, DTU 207, DK-2800 Lyngby, Denmark

⁴ General Physics Institute, Russian Academy of Sciences, 38 Vavilov st., Moscow 117942 Russia

(February 1, 2008)

A free quantum particle in two dimensions with vanishing angular momentum (s-wave) in the form of a ring-shaped wave packet feels an attractive force towards the center of the ring, leading first to an implosion followed by an explosion.

PACS numbers: 03.65.-w, 67.40.Vs, 03.40.Gc

The spreading of a wave packet is an important topic in quantum mechanics [1]. It arises due to different momenta contained in the packet. As pointed out by Max Born [2] even a classical ensemble of particles spreads. The only quantum effect lies in the connection between the position and the momentum distributions, that is, the spreads in position and momentum are not independent. In the present paper we predict a wave effect of the free particle that manifests itself in the dynamics: A ring-shaped wave packet in two dimensions of vanishing angular momentum feels an attractive force towards the center of the ring. This effect is unique to two dimensions and manifests itself most strongly for vanishing angular momentum.

In order to illustrate this surprising phenomenon we first consider the free motion of wave packets in N dimensions. We then concentrate on two and three dimensions, where we propagate a circular shell and a spherical shell, respectively. All wave packets have initially vanishing angular momentum and vanishing average radial momentum. We show that the circular shell first implodes and then explodes. In contrast, the spherical shell does not implode but only explodes. We demonstrate this wave effect unique to two-dimensional space using (i) analytical expressions for the average radial position and momentum, (ii) the Wigner function approach [3] towards quantum mechanics, (iii) analytical expressions [4] for the radial Green's function in two dimensions and (iv) the numerical propagation of wave packets.

Our analytical approach considers the propagation of the initial wave function

$$\Phi_0^{(N)}(x_1, \dots, x_N) = \frac{\mathcal{N}}{\sqrt{S^{(N)}}} r^\gamma \exp \left[-\frac{1}{2} \left(\frac{r}{\delta r} \right)^2 \right] \quad (1)$$

in the absence of any potential. Here $\mathcal{N} \equiv [2/\Gamma(N/2 + \gamma)]^{1/2} \delta r^{-N/2-\gamma}$ denotes the radial normalization constant and $r \equiv (x_1^2 + x_2^2 + \dots + x_N^2)^{1/2}$ is the radial variable in N space dimensions. Moreover, $S^{(N)} \equiv 2\pi^{N/2}/\Gamma(N/2)$ is the total solid angle. The quantity δr defines a characteristic length scale of the packet and is a measure of the radial width [5].

We first take the exponent γ to be equal to 2. This special form of the initial wave function – polynomial in coordinates multiplied by a Gaussian – allows us to calculate analytically the time dependence of the average radial position $\langle r^{(N)} \rangle$ and the average radial momentum $\langle p^{(N)} \rangle$ in N dimensions. We then turn to the general case of arbitrary $\gamma > 1$ but concentrate on two dimensions.

In the derivation of our results we follow three independent approaches that provide identical results: (i) We consider the time evolution in Wigner phase space and calculate the averages with the help of the Wigner function; (ii) we use the method of the generating function [6] to propagate a modulated Gaussian wave packet of the free particle in N dimensions, and evaluate the averages; (iii) we propagate the free wave packet with the help of the radial Green's function of the free particle in two dimensions. Here we do not present the detailed calculations but only summarize the results [7].

We start by presenting first the general results for the propagation in N dimensions and then focus on the special case of two and three dimensions. It is convenient to introduce the dimensionless time variable $\tau \equiv \hbar t / (\delta r^2 M)$, where M denotes the mass of the particle. We find [7] the expression

$$\langle r^{(N)} \rangle(\tau) = \frac{1 + \tau^2/a^{(N)}}{\sqrt{1 + \tau^2}} r_0^{(N)} \quad (2)$$

for the average radial position. Here we have introduced the abbreviation

$$a^{(N)} \equiv 1 + \frac{4N}{N^2 + 3} \quad (3)$$

and the initial radial position

$$r_0^{(N)} \equiv \frac{\Gamma(\frac{N+5}{2})}{\Gamma(\frac{N+4}{2})} \delta r, \quad (4)$$

in N dimensions, where Γ denotes the Euler gamma function.

Equation (2) shows that for large times the average radial position increases linearly with time, corresponding

to an explosion of the wave packet. This holds true for any number of dimensions.

For short times we can expand the square root in Eq. (2) which yields

$$\langle r^{(N)} \rangle(\tau) \simeq \left[1 + \left(\frac{1}{a^{(N)}} - \frac{1}{2} \right) \tau^2 \right] r_0^{(N)}. \quad (5)$$

An interesting situation occurs when $a^{(N)} > 2$. In this case, the average radius initially decreases, corresponding to an implosion of the wave packet. From the definition Eq. (3) of $a^{(N)}$ we arrive at the implosion condition

$$(N-1)(N-3) < 0. \quad (6)$$

Hence an implosion occurs only for $N = 2$ corresponding to $a^{(2)} = 15/7 = 2 + 1/7$.

The phenomenon of the implosion can also be viewed by considering the time evolution of the average radial momentum

$$\langle p^{(N)} \rangle(\tau) = M \frac{d}{dt} \langle r^{(N)} \rangle = \frac{d}{d\tau} \langle r^{(N)} \rangle \frac{\hbar}{\delta r^2}. \quad (7)$$

determined by the time derivative of the average position $\langle r^{(N)} \rangle$, which yields

$$\langle p^{(N)} \rangle(\tau) = -\frac{\tau}{(1+\tau^2)^{3/2}} \left(a^{(N)} - 2 - \tau^2 \right) p_\infty^{(N)} \quad (8)$$

where

$$p_\infty^{(N)} \equiv \lim_{\tau \rightarrow \infty} \langle p^{(N)} \rangle(\tau) = \frac{1}{a^{(N)}} \frac{\hbar r_0^{(N)}}{\delta r^2}. \quad (9)$$

The radial momentum is negative, corresponding to an implosion, provided $\tau < [a^{(N)} - 2]^{1/2}$. Since in two dimensions $a^{(2)} = 15/7$ this condition reads $\tau < \tau_{\min} \equiv 1/\sqrt{7}$. At that moment the momentum vanishes and the implosion turns into an explosion. The average radial position assumes a minimum $r_{\min} \equiv \langle r^{(2)} \rangle(\tau_{\min})$.

In order to highlight the special property of two dimensions we now compare the dynamics of the wave packet Eq. (1) for $\gamma = 2$ in two and three dimensions. In Fig. 1 we show the time evolution of the average radial position and the average radial momentum in two (solid curves) and three (dashed curves) dimensions based on Eqs. (2) and (8) with $a^{(2)} = 15/7$ and $a^{(3)} = 2$.

For short times the average radius of a ring-shaped wave packet decreases as a function of time. This initial implosion of the circular wave is followed by a final explosion. We emphasize that this phenomenon is not due to an external classical force but occurs for a free particle moving in two dimensions [8]. Obviously the spreading of the ring-shaped wave packet in the radial direction is asymmetric; initially the probability flux towards the center is larger than towards the outside.

In the three-dimensional case, where we propagate a spherical shell, the average position and the average momentum do not display the soft implosion effect of two

dimensions. Indeed, since $a^{(3)} = 2$, the average momentum is always positive and the average position cannot decrease but must increase.

The contraction effect is maximal at $\tau_{\min} = 1/\sqrt{7}$ and according to Eq. (2) the average radial position $\langle r^{(2)} \rangle$ assumes the minimal value

$$r_{\min} \equiv \sqrt{\frac{224}{225}} r_0^{(2)} \simeq 0.9978 r_0^{(2)}. \quad (10)$$

Hence, it is a rather soft implosion.

We note that this result is independent of δr , since there is only one length scale in the problem. When we choose a different form for the packet we can make the effect larger. For example, by replacing r^γ in Eq. (1) by $\sin(r^2/\delta r^2)$, we obtain at $\tau_{\min} \simeq 1.11$

$$r_{\min} \simeq 0.9964 r_0^{(2)}. \quad (11)$$

In order to study the implosion effect we concentrate on two dimensions and now consider an arbitrary value of the exponent $\gamma > 1$ in the wave packet Eq. (1). With the help of the corresponding radial propagator [4] we obtain the radial wave function which yields the expression

$$\langle r^{(2)} \rangle(\tau) = 2^\gamma \frac{\Gamma^2(\frac{\gamma+1}{2})}{\Gamma(\gamma + \frac{1}{2})} \frac{\tau^{\gamma-1}}{(1+\tau^2)^{\frac{\gamma+1}{2}}} I(\tau; \gamma) r_0^{(2)} \quad (12)$$

for the average radial position. Here the integral

$$I(\tau; \gamma) \equiv \int_0^\infty d\xi \xi^2 \exp\left(-\frac{1}{1+\tau^2} \xi^2\right) \left| {}_1F_1\left[-\frac{\gamma-1}{2}; 1; \frac{i}{2\tau} \frac{1}{1+i\tau} \left(\frac{r}{\delta r}\right)^2\right] \right|^2 \quad (13)$$

containing the confluent hypergeometric function ${}_1F_1$ can be expressed in terms of the generalized hypergeometric function ${}_3F_2$. However, we found it more convenient to evaluate this integral numerically. For each value of γ we obtain an implosion but the ratio $r_{\min}/r_0^{(2)}$ is always of the same order of magnitude.

We now turn to the explanation of the implosion effect using the Wigner function [3]. The free particle is one of the few quantum mechanical systems in which the classical phase space dynamics is identical to the quantum mechanical one. Indeed, the Wigner function satisfies the classical Liouville equation. Hence, we find the Wigner function at time t from the Wigner function at time $t = 0$ by replacing the position $\vec{r}$ by $\vec{r} - \vec{p}t/M$. We emphasize that now $\vec{r}$ and $\vec{p}$ denote N -dimensional vectors. The corresponding probability distributions in position and momentum follow by integration over the conjugate variable.

Since for the free particle classical and quantum phase space dynamics are identical the quantum effect of implosion rather than explosion is stored in the initial

Wigner function. To be more precise, it is in the correlations between position and momentum. A classical phase space distribution enjoys a probability interpretation and therefore always has to be positive. In contrast, the Wigner function can assume negative values. Indeed, both the Wigner functions of the circular shell and of the spherical shell contain domains in phase space where they become negative. Since the Wigner function is normalized the total volume of the Wigner function is unity. We can calculate numerically the volumes $V_-^{(2)}$ and $V_-^{(3)}$ corresponding to the negative parts. We find in the two dimensional case $V_-^{(2)} \simeq 0.27$, whereas in three dimension $V_-^{(3)} \simeq 0.23$. Hence, there is a slightly larger contribution of negative parts in two dimensions than in three dimensions.

In the phase space description of quantum mechanics quantum effects are stored in the negative parts of the initial Wigner function. Since a classical ensemble of particles cannot have a negative phase space distribution it cannot display the implosion effect. Needless to say a classical ensemble with a ring-shaped initial position distribution can show an implosion provided there exists a non-vanishing negative initial radial momentum. However, in our quantum mechanical wave packet the average radial momentum vanishes.

For the choice Eq. (1) of the initial wave packet the radial probability density

$$\begin{aligned} W^{(N)}(r)dr &= |\Phi_0^{(N)}(r)|^2 r^{N-1} dr S^{(N)} \\ &= |\mathcal{N}|^2 r^{N+2\gamma-1} e^{-(r/\delta r)^2} dr \end{aligned} \quad (14)$$

depends on the number of dimensions. Hence, one might argue that the implosion is a result of the slightly different initial radial wave functions. In order to exclude this argument we have performed a numerical integration of the Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} u_0^{(N)}(r, t) = \left[-\frac{\hbar^2}{2M} \frac{\partial^2}{\partial r^2} + V^{(N)}(r) \right] u_0^{(N)}(r, t) \quad (15)$$

for the radial wave function $u_0^{(N)}$ of the free particle in two and three space dimensions, corresponding to zero angular momentum. Here we always start with the same initial radial wave function. In this way we guarantee identical radial probabilities densities in two and three dimensions.

The radial potential

$$V^{(N)}(r) \equiv \frac{\hbar^2}{2M} \frac{(N-1)(N-3)}{4r^2} \quad (16)$$

in N dimensions involves the product $(N-1)(N-3)$ familiar from the implosion condition Eq. (6).

In three dimensions the potential $V^{(3)}$ vanishes. However, for $N = 2$ the quantum anti-centrifugal potential

$$V_Q(r) \equiv V^{(2)}(r) \equiv -\frac{\hbar^2}{2M} \frac{1}{4r^2} \quad (17)$$

does not vanish and is negative, that is, attractive [9,10]. This potential is responsible for the implosion.

A large class of wave packets with vanishing angular momentum displays this effect of first imploding and then exploding [11]. Indeed, in the insert of Fig. 1 we show the short time evolution of the average radial position and the average radial momentum in two and three dimensions for an initial radial Gaussian wave packet $u(r, t = 0) \equiv \tilde{\mathcal{N}} \exp \left[-(r - \rho)^2 / \delta r^2 \right]$ displaced by an amount ρ from the origin. As in the analytical results we find an implosion of a circular but not of a spherical shell.

So far we have concentrated on a single particle but the effect of implosion manifests itself also in the relative motion of two free particles. We can envision the total wave function of the two particles to be of the form Eq.(1) corresponding to an entangled state. When we assume that each particle is allowed to move in a D -dimensional space the total wave function lives in a $N = 2D$ dimensional configuration space. The physical situation of a single particle imploding in two dimensions corresponds to two particles constrained to one dimension. When they move along a line their average relative separation first decreases and then increases. It looks like the particles first attract and then repel each other. However, when they move in two or three dimensions, their relative distance always increases.

We conclude by briefly outlining an experiment to demonstrate the implosion effect. At first sight, one might think of cold atoms around a wire familiar from atom chips [12]. However, it might be easier to make use of the analogy between the Schrödinger equation and the paraxial wave equation of classical electrodynamics. We shine monochromatic light onto an opaque screen with an annular aperture. In order to avoid diffraction from sharp edges, we introduce apodization, that is, a position dependence transmission coefficient. At the edges of the ring almost no light is transmitted, whereas at the center of the ring almost all light passes through. This arrangement creates an intensity distribution after the screen confined mainly to a ring. The subsequent free propagation in space is analogous to the free propagation in time of the Schrödinger wave function. The time dependent probability distribution then corresponds to the transverse intensity distributions at different locations downstream from the screen [13].

We thank I. Sh. Averbukh, M. V. Berry, D. Bouwmeester, H. Carmichael, D. Greenberger, D. Kobe, R. F. O'Connell, W. C. Schieve, F. Straub, S. Varro and K. Wódkiewicz for many fruitful discussions. I. B.-B., J. P. D. and M. F. gratefully acknowledge the support of the Humboldt foundation and the great hospitality enjoyed at the Abteilung für Quantenphysik. The work of W. P.

-
- [1] D. Bohm, *Quantum Theory*, (Prentice Hall, Englewoods Cliffs, 1951).
 - [2] M. Born, *Science*, **122**, 675 (1955).
 - [3] For an introduction and overview on the concept of Wigner function, see M. Hillery, R. F. O'Connell, M. O. Scully and E. P. Wigner, *Phys. Rep.* **106**, 121 (1984); J. P. Dahl in *Conceptual Trends in Quantum Chemistry*, E. S. Kryachko and J. L. Calais eds., (Kluwer, The Netherlands, 1994), p.199; W. P. Schleich, *Quantum Optics in Phase Space*, (Wiley-VCH, Berlin, 2001).
 - [4] M. A. Cirone, J. P. Dahl, M. Fedorov, D. Greenberger, and W. P. Schleich, submitted to *J. Phys. B*; preprint quant-ph/0108083.
 - [5] We emphasize that δr is *not* the second moment $\Delta r^2 \equiv \langle r^2 \rangle - \langle r \rangle^2$ of the probability distribution $|\Phi_0^{(N)}|^2$. Indeed, Δr depends on the number N of dimensions, whereas δr is a length scale independent of N .
 - [6] I. Białynicki-Birula, Z. Białynicka-Birula and C. Sliwa, *Phys. Rev. A* **61**, 032110 (2000).
 - [7] I. Białynicki-Birula, M. A. Cirone, J. P. Dahl, M. Fedorov, and W. P. Schleich, in preparation.
 - [8] The phenomenon of an imploding wave packet is reminiscent of the transient effect of a free wave packet contracting rather than spreading when it is initially prepared in a contractive state, see for example D. F. Walls and G. Milburn, *Quantum Optics* (Springer, Heidelberg, 1994) p.337. However, the implosion is a purely wave effect, whereas the contractive state is a purely classical effect. It is due to propagation in phase space starting from a clever choice of the initial wave packet.
 - [9] See for example S. Flügge, *Rechenmethoden der Quantenmechanik* (Springer, Heidelberg, 1965).
 - [10] M. A. Cirone, G. Metikas and W. P. Schleich, *Z. Naturforsch.* **56a**, 48 (2001); M. A. Cirone, K. Rzążewski, W. P. Schleich, F. Straub, and J. A. Wheeler, *Phys. Rev. A* (accepted for publication).
 - [11] A purely Gaussian wave packet located initially at the origin does not implode. It spreads symmetrically in radial direction. From Heisenberg's equations of motion for the radial variable or from Ehrenfest's theorem we can derive conditions on the wave function for the implosion effect to occur, see I. Białynicki-Birula, J. P. Dahl, and W. P. Schleich, to be published.
 - [12] J. Denschlag, D. Cassettari, and J. Schmiedmayer, *Phys. Rev. Lett.* **82**, 2014 (1999); E. A. Hinds, C. J. Vale and M. G. Boshier, *Phys. Rev. Lett.* **86**, 1462 (2001). For a nice summary of magnetic chips and quantum circuits for atoms, see E. A. Hinds, *Physics World* **14** (7), 39 (2001).
 - [13] The implosion effect is not the wave mechanics analogue of the Poisson spot [14] familiar from the diffraction theory of classical optics. The implosion results from the modification factor $\mathcal{H}$ discussed in Ref. [4] in Eq.(15) of the free Green's function. A Poisson spot occurs al-

ready when we approximate the radial propagator in two dimensions by the difference of two one-dimensional Green's functions of free motion. Hence, the implosion effect is a modification on top of the Poisson spot.

- [14] M. Born and E. Wolf, *Principles of Optics* (Cambridge University Press, Cambridge, 1999) p.417.

FIG. 1. Comparison between the time evolutions of a circular (solid line) and a spherical shell (dashed line) using the average radial position (top) and the average radial momentum (bottom) obtained from Eqs.(2) and (8) for $a^{(2)} = 15/7$ and $a^{(3)} = 2$. On average the two-dimensional shell first contracts and then expands, whereas the three-dimensional one expands from the onset. In the left inserts we show the short time behavior of a displaced Gaussian radial wave packet obtained numerically. Here we have used the same Gaussian for an initial radial wave function in two and three dimensions with $\delta r = 0.4$ and $\rho = 1.5$ in arbitrary units. Position is scaled in units of the initial position $r_0^{(N)}$ of the wave packet, and momentum in terms of the final momentum $p_\infty^{(N)}$.

